


Effect of exercise on the inter-rectus distance in pregnant women with diastasis recti abdominis: an experimental longitudinal study

Nina-Margrethe Theodorsen, Rolf Moe-Nilssen, Kari Bø, Inger Haukenes



PII: S0031-9406(23)00051-2

DOI: <https://doi.org/10.1016/j.physio.2023.08.001>

Reference: PHYST1356

To appear in: *Physiotherapy*

Please cite this article as: Nina-Margrethe Theodorsen, Rolf Moe-Nilssen, Kari Bø and Inger Haukenes, Effect of exercise on the inter-rectus distance in pregnant women with diastasis recti abdominis: an experimental longitudinal study, *Physiotherapy*, (2023) doi:<https://doi.org/10.1016/j.physio.2023.08.001>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier.

**Title:**

Effect of exercise on the inter-rectus distance in pregnant women with diastasis recti abdominis: an experimental longitudinal study

**Title:**

Effect of exercise on the inter-rectus distance in pregnant women with diastasis recti abdominis: an experimental longitudinal study

**Authors:**

Nina-Margrethe Theodorsen<sup>a</sup>, PT, MSc

Rolf Moe-Nilssen<sup>a</sup>, PT, PhD

Kari Bø<sup>b</sup>, PT, PhD

Inger Haukenes<sup>a</sup>, PT, PhD

**Author Information:**

<sup>a</sup>University of Bergen, Department of Global Public health and Primary Care, PO. Box 7804, 5020 Bergen, Norway.

<sup>b</sup>Norwegian School of Sport Sciences, Department of Sports Medicine, Postboks 4014 Ullevål stadion, 0806 Oslo, Norway.

**Corresponding author:**

Nina-Margrethe Theodorsen, PhD., University of Bergen, Department of Global Public Health and Primary Care, Postboks 7804, NO-5020 Bergen

[nina-margrethe.theodorsen@uib.no](mailto:nina-margrethe.theodorsen@uib.no)

**Word count:** 3095

## Abstract

**Objective:** To investigate the effect of acute contraction of the pelvic floor muscles (PFM) and abdominal exercises on the inter-rectus distance (IRD) compared to resting values, and differences between gestation weeks 27 and 37, in pregnant women with diastasis recti abdominis (DRA).

**Design:** Experimental longitudinal design.

**Setting:** Physiotherapy clinic, primary health care.

**Participants:** Thirty-eight pregnant women with DRA  $\geq 2.8$  cm.

**Interventions:** Two-dimensional ultrasound images of IRD 2 cm above and below the umbilicus were taken at rest and during PFM and abdominal exercises at gestation week 27 and 37. Repeated measures analyses of variance (ANOVAs) with post hoc tests was performed for each exercise for both locations and timepoints.

**Main Outcome measures:** Change in IRD.

**Results:** There was a mean increase of the IRD from rest during a PFM contraction (2mm, 95% CI: 2, 3), drawing-in (4mm, 95% CI: 3, 5) and a combination of these (5 mm, 95% CI: 4, 6) There was a mean decrease of the IRD from rest during the headlift (-3 mm, 95% CI: -4, -2), the curl-up (-3 mm, 95% CI: -4, -2) and the diagonal curl up (-4 mm, 95% CI: -5, -3). Effect of time from gestation week 27 to 37 was a mean increase of 8 mm (95% CI: 6, 9).

**Conclusion:** Pelvic floor and drawing-in exercise increased the IRD, whilst headlift, curl up and diagonal curl up decreased the IRD in pregnant women with DRA at gestation week 27 and 37.

**Contribution of the paper:**

- This is the first study to investigate the acute effect of exercise on IRD in pregnant women with DRA.
- Diagonal curl-up, headlift and curl-up reduced the IRD in pregnant women with DRA at both gestation week 27 and 37. The drawing-in exercise and a PFM contraction increased the IRD at the same timepoints.
- As exercise involving the rectus abdominis muscle decrease the inter-rectus distance in pregnant women, clinicians should reconsider advising against such exercises.

**Key words:** *diastasis, exercise, linea alba, pregnancy, ultrasonography*

## Introduction

Diastasis recti abdominis (DRA) is a common condition in pregnant and postpartum women, with a reported prevalence of up to 100% during the last trimester of pregnancy, 60% 6 weeks postpartum and 32% 12 months postpartum (1, 2). DRA is defined as a stretching and weakness of the Linea alba increasing the distance between the two muscle bellies of the rectus abdominis muscle (3). During pregnancy, this stretching and weakness is a result of the increased strain on the ventral abdominal wall caused by the growing foetus (3, 4, 5). DRA is diagnosed by measuring the distance between the medial borders of the two rectus abdominis muscle bellies, referred to as the inter-rectus distance (IRD). There is however, no consensus on either the cut-off point for a clinically significant DRA, or where to measure the distance along the Linea alba (6). Evidence suggest that DRA is associated with weak abdominal muscles and increased abdominal pain (7). DRA is also believed to be associated with pelvic girdle pain, low back pain and urinary incontinence, however the evidence for these associations are weak (8, 9, 10).

Both palpation by finger width and calipers are considered valid and reliable screening methods for measuring IRD in clinical settings, however, ultrasound imaging of the IRD is the most reliable and recommended measurement tool when measuring and monitoring change in the IRD (6, 11).

Physiotherapists in primary health care commonly treat women, both during pregnancy and the postpartum period, presenting with DRA (12). Therapy aims for both prevention and reduction of DRA, however current clinical recommendations build on weak or sparse evidence as to which exercise modality is the most effective and feasible (4, 13, 14). Based on well-established clinical beliefs pregnant women are

often advised against exercises activating the rectus abdominis muscle as this is assumed to increase DRA (12, 15).

Most studies on DRA have been conducted on postpartum populations, and to date there are no basic studies investigating the effect of different exercises on the IRD in pregnant women. As DRA mostly occurs and develops during 2<sup>nd</sup> and 3<sup>rd</sup> trimester, it is important to examine how the IRD is influenced by exercise during this period (1, 2). Identifying exercise with a potential influence on the IRD, is the first stage of knowledge in how to reduce the IRD in women with DRA and particularly the unresolved prevalence (32%) 12 months postpartum (2). The aim of this present study was to identify the acute effect of a pelvic floor muscle (PFM) contraction and 4 different abdominal muscle exercises on the IRD in pregnant women presenting with DRA, measured at gestation week 27 and 37. Difference in between assessments were also investigated.

## Methods

### Design

A longitudinal experimental design with repeated measures was applied, where data was collected from all participants at two timepoints; gestation week 27 and gestation week 37.

### Participants

Thirty-eight pregnant women were included, based on a power calculation performed with data from two previous studies where a clinically significant difference of 5 mm was used between IRD 2 cm above the umbilicus at rest and IRD during contraction of transversus abdominis muscle, with an estimated standard deviation

(SD) for the difference of 11 mm (11, 16). A significance level .05 and power of 0.80 was applied.

Participants were recruited from local physiotherapy clinics, local health centres, and through social media platforms Facebook and Instagram.

The inclusion criteria were healthy pregnant primigravida and parous women aged > 18 years, less than 27 weeks pregnant. In accordance with Mota et al (17) the participants had to present with a IRD of 2.8 cm or more, measured 2 cm above and/or below the umbilicus on initial assessment. Studies report the largest change in IRD is measured to 2 cm above the umbilicus, hence measurements were done 2 cm above and 2 cm below the umbilicus. Both cut-off point and measurement locations were set in accordance with similar studies on postpartum populations to ensure reliable measurements and comparisons of results (11, 16, 17, 18). Participants presenting with an abdominal wall protrusion along the linea alba were included, even though the IRD was measured to less than 2.8 cm, as suggested by Brauman et al (2018) (19).

Exclusion criteria were pregnancies where exercise was contraindicated, serious illnesses regarding both mother and/or foetus, inability to understand and speak a Scandinavian language, failure to complete and present an informed consent and chronic physical or mental illness incompatible with the intervention. Exclusion criteria during the study period were stillbirth or premature birth before gestation week 37.

Participants were given oral and written information about the study. The study was approved by the Regional medical Ethics Committee (76296 2020/2304) and procedures agreed with the Helsinki Declaration.

## Primary outcome measure

Change between IRD during exercise and at rest measured in mm (6, 20).

## Instrumentation and Examiner

Ultrasound images (B-mode) of the IRD were collected using a two-dimensional ultrasound diagnostic scanner with a linear probe (Alpinion Ecube8). Ultrasound imaging is found to be a reliable measurement method for the IRD, with a very good intra-tester reliability (11).

All assessments, images and measurements were performed by one investigator, a women's health physiotherapist with 22 years of clinical experience who had undergone specific training in ultrasound imaging of the pelvic floor and abdomen. Blinding of the investigator was ensured by converting all images to a digital format, allocating them a random number prior to measurements.

## Procedures

All participants completed an online questionnaire on background variables prior to data collection. Age, body mass index, parity, single/multiple fetuses, protrusion, education, and smoking were considered relevant variables to describe the population included. Assessments were performed when the participants were in gestation week 27 and 37. Presence and size of DRA, and the ability to perform a correct PFM contraction and correct abdominal exercises were assessed prior to inclusion in the study. DRA was assessed and measured using the ultrasound machine's integrated measurement tool. Presence of a protrusion was assessed by observation. The participant was instructed to perform an abdominal curl-up and if a doming along the linea alba occurred, the participant was eligible for inclusion. The ability to perform a correct pelvic floor contraction and the drawing-in exercise was assessed by



ultrasonography and are described further, including descriptions of the different exercises, in **TABLE 1**.

The position of the participant during imaging was supine, knees hip-width apart and bent 90°, feet resting on plinth, and the head resting on a pillow. Arms were placed alongside the body or behind the head according to which exercise was performed. Measurement locations were initially marked on the skin to standardize the position of the ultrasound transducer when imaging the IRD; 2 cm above the umbilicus, measured from the upper edge of the umbilicus, and 2 cm below the umbilicus, with the lower edge of the umbilicus as point of reference.

The transducer was placed transversely along the midline of the abdomen and the bottom edge of the transducer corresponded with the skin markers. The transducer was moved laterally until the medial borders of the rectus abdominis muscles, and the Linea alba were visualized. To enable accurate measurements when the IRD was too large to be captured with standardized brightness mode (B-imaging), images were taken using an extended field of view or panoramic function to produce one long image displaying the IRD (21).

The images were collected in a set order. Prior to each imaging the participants were told to relax, take a deep breath in, and exhale before the exercises were performed. Images were collected immediately at the end of exhalation and care was taken so the pressure of the probe did not cause a reflexive response of the abdominal muscles (22). The participants were told to hold the contractions for approximately three to five seconds to allow imaging. One image was collected at each location for each task. Ten images of the IRD at two different locations (2 cm above and below the umbilicus) were collected for each participant: (1) at rest (2) during a PFM contraction, (3) during a drawing-in exercise, (4) during a combined PFM contraction and drawing-

in exercise, (5) during a head lift, (6) during a curl-up, (7) during a combined drawing-in exercise and curl-up, (8) during a combined PFM contraction, a drawing-in exercise and curl-up, (9) during a diagonal curl-up to the left, and (10) during a diagonal curl-up to the right.

Ultrasound images were stored on the ultrasound machine as Digital Imaging and Communications in Medicine (DICOM) format. The images were then transferred and stored at the SAFE server, University of Bergen, the University's information technology department database. Here, all participants images were allocated a random number to ensure blinding of the assessor. Measurements of the IRD were done using the MATLAB Image Processing Toolbox programme. The medial borders of rectus abdominis (outside the hyperechoic line) and the conjunction to the Linea alba on both sides were identified and the transverse linear distance between these borders, were measured. This measurement procedure is in line with the procedure described by Mota et al (11).

### Statistical Analysis

SPSS (Version 28; IBM Corporation) was used for all statistical analyses. Background variables are presented as means with standard deviation and ranges, or number and percentages. Measures of IRD were presented as means with standard deviation and tested for normality using the Shapiro-Wilks test. Repeated-measures analysis of variance (ANOVA) with post hoc tests was performed for each exercise (above and below the umbilicus, and gestation week 27 and 37). Results were presented as means with 95% confidence interval (CI); statistical significance was set at  $P < .05$  with Bonferroni correction for multiple comparisons. 95% Limits of Agreement in mean IRD between test and retest has previously been reported to be between -8.7 and 8.3 mm (11).

The effect of each exercise was defined as change in IRD from rest to each contraction, and the results are presented with a positive value with an increase in IRD during the exercise and a negative value with a decrease in IRD during the exercise. The diagonal curl-up values are presented as an average between left and right values.

## Results

Thirty-eight pregnant women completed both assessments, and demographic data are presented in **TABLE 2**. The descriptive statistics for the IRD measurements at gestation weeks 27 and 37 are presented in **TABLE 3** and **TABLE 4**.

The variable of interest was the change in IRD between rest and 9 different exercises. The post hoc analysis indicates a mean increase in IRD during a PFM contraction of 2 mm (95% CI: 2, 3),  $p < .001$ , during a drawing-in exercise of 4 mm (95% CI: 3, 5),  $p < .001$ , and during a combined PFM contraction and the drawing-in exercise of 5mm (95% CI: 4, 6),  $p < .001$ . The post hoc analysis indicates a mean decrease in IRD during a headlift of -3 mm (95% CI: -4, -2),  $p < .001$ , during a curl-up of -3 mm (95% CI: -4, -2),  $p < .001$ , and during a diagonal curl-up of -4mm (95% CI: -5, -3),  $p < .001$ . The combined exercises drawing-in with curl-up and PFM contraction with drawing-in and curl-up showed no significant change from the resting value (Figure 1).

These findings showed a main effect of location with 4 mm higher values above compared to below the umbilicus (95% CI 3, 5),  $p < .001$ , and a main effect of time with an increase from gestation week 27 to 37 of 8 mm (95% CI: 6, 9),  $p < .001$ , (Figure 1).

## Discussion

Overall, this study demonstrates that the instantaneous impact of exercise on the IRD in pregnant women with DRA varies by the exercises performed. These results

were consistent both above and below the umbilicus, and at gestation week 27 and 37. The results can be divided into three main trends, where (1) the IRD increases significantly with pelvic floor and transversus abdominis activation, (2) decreases significantly with headlift, curl-up and diagonal curl-up, and (3) shows no significant change compared to rest where exercises from the two groups are combined.

#### Exercises increasing the IRD

The present study found that the IRD increased significantly with a PFM contraction. These findings are in line with studies performed on postpartum populations using equivalent measurement methods, and hence adds strength to the view that PFM has no specific role in reducing the IRD (10, 13, 16, 18, 23). Pregnant women are encouraged to do PFM training throughout the pregnancy and postpartum period, and there is Level 1 evidence (recommendation A) that PFM training should be the first treatment choice to prevent and reduce urinary incontinence (UI) (24). Woodley et al (24) found that continent pregnant women have 62% less risk of developing UI in late pregnancy and 29% during the mid-postnatal period if they do PFM training (24).

Even though we found a significant increase in IRD during a PFM contraction, this change was minimal. The longitudinal effect of PFM training on the IRD is yet to be investigated, as this study can only determine an immediate effect due to the study design. Pregnant women should still be encouraged to perform regular PFM training, but clinicians should be cautious postulating synergistic effects between the PFM and reduction of the IRD.

The largest increase of the IRD in this study, was measured during a combined PFM contraction and a drawing-in exercise activating the transversus abdominis. There is a synergistic co-contraction between the PFM and the transversus abdominis, so that when the PFM contracts, it enhances the contraction of the transversus

abdominis (25, 26). This supports our findings as a combination of a PFM contraction and drawing-in exercise increased the IRD significantly more than an isolated drawing-in exercise. Similar effects have been reported in studies investigating the effect of pelvic floor and transversus abdominis muscle contraction on IRD in a postpartum population with DRA (16, 18).

The activation of the transversus abdominis through a drawing-in exercise is postulated to be the gentlest and most effective to restore form and function of the linea alba (4, 27). Contraction of the transversus abdominis was previously thought to reduce the IRD, but recent studies have contradicted this assumption in the postpartum and general female population (18, 28). Our findings in a pregnant population are also in line with this, as the IRD significantly increased during a drawing-in exercise. This contrasts with common physiotherapy practice, as Keeler et al (12) report that most women health physiotherapists prescribe transversus abdominis (89%) and pelvic floor (87%) exercise to rehabilitate DRA (12). One study has suggested that the aim of treatment ought to shift from reduction of IRD to function of the linea alba, and that transversus abdominis play a key role in restoring function despite increasing the IRD (28). However, the measurement methods they applied have not been validated or tested for reliability and there is no evidence for this new theory. Literature investigating effects of surgical repairs of DRA, describes reducing the IRD as the main aim by restoring the rectus abdominis to the midline. This will both increase the function of the abdominal wall and enhances the patient's quality of life (29, 30).

#### Exercises decreasing the IRD

Our findings show that diagonal curl-ups decrease the IRD the most. The curl-up and the headlift also reduced the IRD significantly, all in line with recent experimental studies investigating the effect of exercises involving the rectus abdominis on the IRD

in postpartum women with DRA (16, 18, 31). These findings contrast with the advice given to pregnant women not to do exercise involving the rectus abdominis in the belief that this may add pressure to an already stretched linea alba (8, 19) and further increase the IRD (12, 27, 28). There is no evidence supporting this advice and based on the results of the present study clinicians should reconsider advice given to pregnant women regarding abdominal exercise. Randomised clinical trials on a pregnant population are warranted to further investigate whether exercise of the rectus abdominis during pregnancy can prevent or reduce presence of DRA during the postpartum period.

#### Combined exercises

The drawing-in exercise significantly increased the IRD, while the curl-up significantly decreased the IRD. When combined, the IRD did not change significantly either way from the resting values. It may seem that when combined, the two exercises counteract each other's immediate effect on the IRD. This was also reported in a systematic review investigating the evidence for abdominal exercise in postpartum population with DRA (13). This suggest that curl-ups do not have the postulated negative effect on the IRD, and that curl-ups and exercise involving rectus abdominis may play a more significant role in reducing the IRD in patients with DRA. It may also indicate that normal movement involving both layers of abdominal musculature will not increase the IRD in women with DRA.

#### Strengths and weaknesses

To our knowledge this is the first study to investigate the effect of exercise on the IRD in a pregnant population. Strengths of this study are the number of participants based on an apriori power calculation and the longitudinal design. Measurements at two time points during pregnancy increase the power as the results are consistent

between gestation week 27 and 37. Other strengths are blinding of the investigator, use of ultrasound for assessments of the IRD, measurements of 9 different exercises and measurements at both above and below the umbilicus. Participants included in this study were both primi- and multigravida resulting in a sample more reflective of the general population and hence increasing the external validity of the findings. Cut-off point for DRA, standardized assessments and imaging procedures were in accordance with similar studies done on both pregnant and postpartum populations to ensure comparison of results (16, 18, 31).

As there were 20 images for each participant at each timepoint we chose to do the exercises in a set order, to ensure that all images were taken. This might have caused an order effect, and hence, be a limitation of the study. However, we concluded that it was more important to obtain all images, than to counteract a potential order effect caused by fatigue among the participants. Ethnicity was not registered, and all participants were self-sampled. Both these factors may jeopardize representation of the general pregnant population. Another limitation is the study design, in which cross sectional data was analysed. This excludes the ability to conclude whether the effects of different exercises on the IRD are maintained over time.

## Conclusion

The IRD in pregnant women increases with PFM and transversus abdominis exercise, and decreases with a headlift, curl-up and diagonal curl-up, both above and below the umbilicus, and at gestation week 27 and 37. This present study adds valuable novel knowledge of how the IRD in pregnant women with DRA is affected by different abdominal exercises.

**Ethical approval:**

The study was approved by the Norwegian Regional Medical Ethics Committee, with ethical approval number 76296 2020/2304.

**Funding statement:**

This work was supported by The Norwegian Fund for Post-Graduate Training in Physiotherapy, grant number 129673.

**Statement of financial disclosure and conflict of interest:**

None declared.

**Authors` Contributions statement:**

Each author of this paper meets the criteria for authorship. NMT, KB, IH and RMN designed the study. NMT drafted the article and RMN, IH and KB critically revised the article for important intellectual content. The final manuscript was seen and approved by all authors.

**Data sharing statement:**

There are no data available.



1. Mota PGFD, Pascoal AGBA, Carita AIAD, Bø K. Prevalence and risk factors of diastasis recti abdominis from late pregnancy to 6 months postpartum, and relationship with lumbo-pelvic pain. *Manual Therapy*. 2015;20(1):200-5.
2. Sperstad JB, Tennfjord MK, Hilde G, Ellström-Engh M, Bø K. Diastasis recti abdominis during pregnancy and 12 months after childbirth: prevalence, risk factors and report of lumbopelvic pain. *British Journal of Sports Medicine*. 2016;50(17):1092.
3. Akram J, Matzen SH. Rectus abdominis diastasis. *J Plast Surg Hand Surg*. 2014;48(3):163-9.
4. Benjamin DR, van de Water ATM, Peiris CL. Effects of exercise on diastasis of the rectus abdominis muscle in the antenatal and postnatal periods: a systematic review. *Physiotherapy*. 2014;100(1):1-8.
5. Boissonnault JS, Blaschak MJ. Incidence of diastasis recti abdominis during the childbearing year. *Phys Ther*. 1988;68(0031-9023 (Print)):1082-6.
6. van de Water ATM, Benjamin DR. Measurement methods to assess diastasis of the rectus abdominis muscle (DRAM): A systematic review of their measurement properties and meta-analytic reliability generalisation. *Manual Therapy*. 2016;21:41-53.
7. Gluppe S, Ellström Engh M, Kari B. Women with diastasis recti abdominis might have weaker abdominal muscles and more abdominal pain, but no higher prevalence of pelvic floor disorders, low back and pelvic girdle pain than women without diastasis recti abdominis. *Physiotherapy*. 2021;111:57-65.
8. Cavalli M, Aiolfi A, Bruni PG, Manfredini L, Lombardo F, Bonfanti MT, et al. Prevalence and risk factors for diastasis recti abdominis: a review and proposal of a new anatomical variation. *Hernia*. 2021;25(4):883-90.
9. Sokunbi G, Camino-Willhuber G, Paschal PK, Olufade O, Hussain FS, Shue J, et al. Is Diastasis Recti Abdominis Associated With Low Back Pain? A Systematic Review. *World Neurosurgery*. 2023.
10. Bø K, Hilde G, Tennfjord MK, Sperstad JB, Engh ME. Pelvic floor muscle function, pelvic floor dysfunction and diastasis recti abdominis: Prospective cohort study. *Neurourology and Urodynamics*. 2017;36(3):716-21.
11. Mota P, Pascoal AG, Sancho F, Bø K. Test-retest and intrarater reliability of 2-dimensional ultrasound measurements of distance between rectus abdominis in women. *The Journal of orthopaedic and sports physical therapy*. 2012;42(11):940.
12. Keeler J, Albrecht M, Eberhardt L, Horn L, Donnelly C, Lowe D. Diastasis Recti Abdominis: A Survey of Women's Health Specialists for Current Physical Therapy Clinical Practice for Postpartum Women. *Journal of Women's Health Physical Therapy*. 2012;36(3):131-42.
13. Gluppe S, Engh ME, Bø K. What is the evidence for abdominal and pelvic floor muscle training to treat diastasis recti abdominis postpartum? A systematic review with meta-analysis. *Brazilian Journal of Physical Therapy*. 2021.
14. Benjamin DR, Frawley HC, Shields N, Peiris CL, van de Water ATM, Bruder AM, et al. Conservative interventions may have little effect on reducing diastasis of the rectus abdominis in postnatal women – A systematic review and meta-analysis. *Physiotherapy*. 2023;119:54-71.
15. Dufour S, Bernard S, Murray-Davis B, Graham N. Establishing Expert-Based Recommendations for the Conservative Management of Pregnancy-Related Diastasis Rectus Abdominis: A Delphi Consensus Study. *Journal of Women's Health Physical Therapy*. 2019;43(2):73.
16. Theodorsen NM, Strand LI, Bø K. Effect of pelvic floor and transversus abdominis muscle contraction on inter-rectus distance in postpartum women: a cross-sectional experimental study. *Physiotherapy*. 2019;105(3):315-20.
17. Mota P, Pascoal AG, Carita AI, Bø K. Normal width of the inter-recti distance in pregnant and postpartum primiparous women. *Musculoskeletal Science and Practice*. 2018;35:34-7.
18. Gluppe SB, Engh ME, Bø K. Immediate Effect of Abdominal and Pelvic Floor Muscle Exercises on Interrecti Distance in Women With Diastasis Recti Abdominis Who Were Parous. (1538-6724 (Electronic)).
19. Brauman D. Diastasis Recti: Clinical Anatomy. *Plastic and Reconstructive Surgery*. 2008;122(5):1564-9.

20. Mota P, Pascoal AG, Sancho F, Carita AI, Bø K. Reliability of the inter-rectus distance measured by palpation. Comparison of palpation and ultrasound measurements. *Manual Therapy*. 2013;18(4):294-8.
21. Keshwani N, Mathur S, McLean L. Validity of Inter-rectus Distance Measurement in Postpartum Women Using Extended Field-of-View Ultrasound Imaging Techniques. *J Orthop Sports Phys Ther*. 2015;45(10):808-13.
22. Teyhen DS, Miltenberger CE, Deiters HM, Del Toro YM, Pulliam JN, Childs JD, et al. The use of ultrasound imaging of the abdominal drawing-in maneuver in subjects with low back pain. *The Journal of orthopaedic and sports physical therapy*. 2005;35(6):346.
23. Wang Q, Yu X, Chen G, Sun X, Wang J. Does diastasis recti abdominis weaken pelvic floor function? A cross-sectional study. *International Urogynecology Journal*. 2019:1-7.
24. Woodley SJ, Lawrenson P, Boyle R, Cody JD, Mørkved S, Kernohan A, et al. Pelvic floor muscle training for preventing and treating urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database of Systematic Reviews*. 2020(5).
25. Bø K, Sherburn M, Allen T. Transabdominal ultrasound measurement of pelvic floor muscle activity when activated directly or via a transversus abdominis muscle contraction. *Neurourology and Urodynamics*. 2003;22(6):582-8.
26. Pereira LC, Botelho S, Marques J, Amorim CF, Lanza AH, Palma P, et al. Are transversus abdominis/oblique internal and pelvic floor muscles coactivated during pregnancy and postpartum? *Neurourology and Urodynamics*. 2013;32(5):416-9.
27. Lee DG, Lee LJ, McLaughlin L. Stability, continence and breathing: The role of fascia following pregnancy and delivery. *Journal of Bodywork & Movement Therapies*. 2008;12(4):333-48.
28. Lee D, Hodges PW. Behavior of the Linea Alba During a Curl-up Task in Diastasis Rectus Abdominis: An Observational Study. *The Journal of orthopaedic and sports physical therapy*. 2016;46(7):580.
29. Mommers E, Ponten J, Al Omar A, Vries Reilingh T, Bouvy N, Nienhuijs S. The general surgeon's perspective of rectus diastasis. A systematic review of treatment options. *And Other Interventional Techniques Official Journal of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) and European Association for Endoscopic Surgery (EAES)*. 2017;31(12):4934-49.
30. Emanuelsson P, Gunnarsson U, Dahlstrand U, Strigård K, Stark B. Operative correction of abdominal rectus diastasis (ARD) reduces pain and improves abdominal wall muscle strength: A randomized, prospective trial comparing retromuscular mesh repair to double-row, self-retaining sutures. *Surgery*. 2016;160(5):1367-75.
31. Mota P, Pascoal AG, Carita AI, Bø K. The Immediate Effects on Inter-rectus Distance of Abdominal Crunch and Drawing-in Exercises During Pregnancy and the Postpartum Period. *The Journal of orthopaedic and sports physical therapy*. 2015;45(10):781.

**TABLE 1.**

Operational description of the exercises, including description of how a correct performance was ensured.

Exercise	Operational description	Assessment
<b>PFM contraction</b>	<b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow, arms resting alongside the body <b>Instruction:</b> Squeeze muscles around all pelvic openings, lift upwards and in-wards	Ultrasound imaging using a curved linear array transducer placed in the mid-sagittal plane immediately suprapubically observing the cranioventral displacement of the pelvic floor in a sagittal plane
<b>Drawing-in</b>	<b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow, arms resting alongside the body <b>Instruction:</b> Inhale, and at end of exhalation the umbilicus is drawn down towards the spine, activating the transversus abdominis muscle	Ultrasound imaging of the Transversus abdominis muscle on the lateral abdominal wall at the level of the umbilicus. A correct performed exercise involved minimal activation of the adjacent abdominal muscles.
<b>PFM + drawing-in</b>	<b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow, arms resting alongside the body <b>Instruction:</b> Contract the PFM first, and then add the drawing-in exercise	Observation that exercise was performed in accordance with operational description
<b>Head lift</b>	<b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow, arms resting alongside the body <b>Instruction:</b> Lift and clear the head off the plinth. Shoulders are to stay on the plinth during the manoeuvre.	Observation that exercise was performed in accordance with operational description
<b>Curl-up</b>	<b>Position:</b> Supine, knees hip-width apart and bent 90° feet resting on plinth, head resting on a pillow, both hands behind head <b>Instruction:</b> Lift head and shoulders upwards until the shoulder blades clear the plinth	Observation that exercise was performed in accordance with operational description
<b>Drawing-in + curl-up</b>	<b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow. Both hands behind head	Observation that exercise was performed in accordance with operational description

<b>PFM + drawing-in + curl up</b>	<p><b>Instruction:</b> Start with the drawing-in exercise, and then do a curl-up</p> <p><b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow Both hands behind head</p> <p><b>Instruction:</b> Start with the PFM contraction, then add the drawing-in exercise, and then do a curl-up</p>	Observation that exercise was performed in accordance with operational description
<b>Diagonal curl-up left and right</b>	<p><b>Position:</b> Supine, knees hip-width apart and bent 90°, feet resting on plinth, head resting on a pillow</p> <p><b>Instruction:</b> Placed right arm behind the head, elbow pointing laterally. Keep hand under head and move the elbow towards the left hip, lifting head and clearing the right shoulder blade of the plinth. Repeat on other side.</p>	Observation that exercise was performed in accordance with operational description

---

Abbreviations: PFM, Pelvic Floor Muscle.

**TABLE 2.** Descriptives of the study population (n=38)

	<i>n</i> (%)	Mean (SD)	Range
Age (years)		34 (3)	28-41
Body mass index			
- gestation week 27		27 (5)	20-37
- gestation week 37		28 (5)	20-39
Inter-rectus distance (mm)			
- gestation week 27, au		31 (6)	13-56
- gestation week 37, au		40 (6)	24-58
- gestation week 27, bu		27 (7)	13-56
- gestation week 37, bu		34 (6)	24-58
Parity		0.8 (0.7)	0-2
Single pregnancies	38 (100)		
Protrusion	15 (40)		
University education	34 (90)		
Smoking	0 (0)		

Abbreviations: SD, standard deviation; au, above umbilicus; bu, below umbilicus.

**TABLE 3.** Mean difference in IRD (mm) between exercise and rest among 38 pregnant women in gestation week 27

Measurement location	Exercise versus rest	Mean difference	SD	95% CI	P-value
2 cm above umbilicus	Pelvic floor muscle	2	2	2, 3	<.001
	Drawing-in	4	2	3, 4	<.001
	Pelvic floor + drawing in	5	3	4, 6	<.001
	Headlift*	-3	2	-3, -2	<.001
	Curl-up*	-3	3	-4, -2	<.001
	Drawing-in + curl-up*	-1	3	-2, -0	0.04
	Pelvic floor+drawing-in+curl-up*	0	4	-1, 1	0.96
	Diagonal curl up*	-4	3	-5, -3	<.001
2 cm below umbilicus	Pelvic floor muscle	2	1	2, 2	<.001
	Drawing-in	4	2	3, 5	0.99
	Pelvic floor + drawing in	5	2	4, 6	<.001
	Headlift*	-3	3	-4, -2	<.001
	Curl-up*	-2	3	-3, -1	0.02
	Drawing-in + curl-up*	-0	3	-1, 1	0.00
	Pelvic floor+drawing-in+curl-up*	-1	3	-0, 2	0.13
	Diagonal curl up*	-3	3	-4, -2	<.001

Abbreviations: IRD, inter-rectus distance; SD, standard deviation; CI, confidence interval.

\*Negative value represents a decrease in IRD compared to rest


**TABLE 4.** Mean difference in IRD (mm) between exercise and rest among 38 pregnant women in gestation week 37

Measurement location	Exercise versus rest	Mean difference	SD	95% CI	P-value
2 cm above umbilicus	Pelvic floor muscle	3	2	2, 3	<.001
	Drawing-in	4	2	4, 5	<.001
	Pelvic floor + drawing in	5	2	5, 6	<.001
	Headlift*	-4	3	-5, -3	<.001
	Curl-up*	-4	3	-5, -3	<.001
	Drawing-in + curl-up*	-2	3	-3, -1	<.001
	Pelvic floor+drawing-in+curl-up*	-1	4	-2, 0	0.06
	Diagonal curl up*	-6	3	-7, -5	<.001
2 cm below umbilicus	Pelvic floor muscle	2	2	2, 3	<.001
	Drawing-in	4	2	3, 5	<.001
	Pelvic floor + drawing in	5	3	4, 6	<.001
	Headlift*	-3	3	-4, -2	<.001
	Curl-up*	-2	3	-3, -1	<.001
	Drawing-in + curl-up*	-0	3	-1, 1	0.67
	Pelvic floor+drawing-in+curl-up	1	3	0, 2	0.04
	Diagonal curl up*	-3	3	-4, -2	<.001

Abbreviations: IRD, inter-rectus distance; SD, standard deviation; CI, confidence interval.

\*Negative value represents a decrease in IRD compared to rest

**FIGURE 1.** Mean difference in IRD between exercise and rest measured at 27 and 37 weeks and analyzed with repeated measures ANOVA with post-hoc tests, Bonferroni correction for multiple comparisons



Abbreviations: IRD, inter-rectus distance; ANOVA, analysis of variance; mm, millimeters; CI, confidence interval; pfm, pelvic floor muscle; di, drawing-in; pfm+di, pelvic floor muscle and drawing-in; cu, curl-up; di+cu, drawing-in and curl-up; pfm+di+cu, pelvic floor muscle, drawing-in and curl-up; diag cu, diagonal curl-up.